

SEMIDIRECT DECOMPOSITIONS OF DIAGONAL SEMIDIRECT PRODUCTS AND APPLICATIONS FOR DIHEDRAL GROUPS

PETERIS DAUGULIS*

Abstract. Nonuniqueness of semidirect decompositions of groups is an insufficiently studied question in contrast to direct decompositions. We obtain a characterization of semidirect decompositions for a special case of semidirect product when the twisting homomorphism acts diagonally on a direct product. We give applications of this result in the case of generalized dihedral groups and classic dihedral groups D_{2n} .

Key words. semidirect product, diagonal action, generalized dihedral group

AMS subject classifications. 20E22, 20D40.

1. Introduction.

1.1. Background. The aim of this article is to study semidirect decompositions of groups both in general and special cases.

By the well known Krull-Remak-Schmidt theorem the multiset of isomorphism types of indecomposable direct factors for groups satisfying ascending and descending chain conditions on normal subgroups does not depend on the order of factors. Thus direct decompositions of such groups, e.g. finite groups, may be considered understood.

Few results of this type are known for semidirect and Zappa-Szep decompositions. One can mention the Schur-Zassenhaus theorem as an example.

We consider the case when the base group is a direct product. We present a general result which allows to characterize some semidirect decompositions in the case when the base group is a direct product and the twisting homomorphism acts diagonally. We give applications of this result in the case which is relatively easy to understand - finite dihedral groups, both classic and generalized.

All groups in this article are assumed finite. We use traditional multiplicative notation for general groups and additive notation for abelian groups. In this article the dihedral group of order m is denoted by D_m . The cyclic group of order m is denoted by \mathbb{Z}_m , in additive notation we assume that $\mathbb{Z}_m = \langle 1 \rangle$ and its elements are

*Department of Mathematics, Daugavpils University, Daugavpils, LV-5400, Latvia (peteris.daugulis@du.lv).

residues mod m . For any $m|n$ we usually identify \mathbb{Z}_m with the corresponding subgroup of \mathbb{Z}_n .

1.2. Basic facts about semidirect products.

We remind the reader that an external semidirect product of groups N (*base group*) and K (*extending group*) is the group $N \rtimes_{\varphi} H = (N \times H, \cdot)$ where the group product is defined on the Cartesian product $N \times H$ using a group homomorphism (*twisting homomorphism*) $\varphi \in \text{Hom}(H, \text{Aut}(N))$ as follows: $(n_1, h_1) \cdot (n_2, h_2) = (n_1 \varphi(h_1)(n_2), h_1 h_2)$. Sets $\tilde{N} = N \times \{e_H\}$ and $\tilde{H} = \{e_N\} \times H$ are subgroups in $N \times H$.

A group G is an internal semidirect product of its subgroups N and H if N is a normal subgroup, $G = NH$ and $N \cap H = \{e\}$. If a group G is finite then for G to be an internal semidirect product NH is equivalent to 1) N being normal in G , 2) $|H| \cdot |K| = |G|$ and 3) $N \cap K = \{e\}$. In the internal case the twisting homomorphism $H \rightarrow \text{Aut}(N)$ is given by the map $h \mapsto (n \rightarrow hnh^{-1})$, for any $n \in N$, $h \in H$.

Both expressions will be called semidirect decompositions of G . If the twisting homomorphism is not discussed, we omit it and use the notation \rtimes .

We consider direct product to be a special case of semidirect product with the twisting homomorphism being trivial.

For relevant treatment see [4], [5].

2. Main results.

2.1. Diagonal semidirect products.

2.1.1. Automorphisms of direct products. We introduce a linear algebra style notation for direct products of groups.

Let $G = G_1 \times G_2$. Encode the element (g_1, g_2) as a column $\begin{bmatrix} g_1 \\ g_2 \end{bmatrix}$. If $\varphi \in \text{Aut}(G)$, then $\varphi\left(\begin{bmatrix} g_1 \\ g_2 \end{bmatrix}\right) = \begin{bmatrix} \varphi(g_1, g_2) \\ \varphi(g_1, g_2) \end{bmatrix}$. One can check, that for all relevant parameter values φ_i satisfy the following properties:

1. $\varphi_i(ab, e) = \varphi_i(a, e)\varphi_i(b, e)$,
2. $\varphi_i(e, ab) = \varphi_i(e, a)\varphi_i(e, b)$,
3. $\varphi_i(a, b) = \varphi_i(a, e)\varphi_i(e, b) = \varphi_i(e, b)\varphi_i(a, e)$,

Define $\varphi_{11}(g_1) = \varphi_1(g_1, e)$, $\varphi_{12}(g_2) = \varphi_1(e, g_2)$, $\varphi_{21}(g_1) = \varphi_2(g_1, e)$, $\varphi_{22}(g_2) =$

$\varphi_2(e, g_2)$, for all $g_i \in G_i$. All functions φ_{ij} are group homomorphisms. Thus $\varphi_i(g_1, g_2) = \varphi_i(g_1, e)\varphi_i(e, g_2) = \varphi_{i1}(g_1)\varphi_{i2}(g_2)$.

We can encode action of φ as follows: $\varphi\left(\begin{bmatrix} g_1 \\ g_2 \end{bmatrix}\right) = \left[\begin{array}{c|c} \varphi_{11}(g_1) & \varphi_{12}(g_2) \\ \hline \varphi_{21}(g_1) & \varphi_{22}(g_2) \end{array}\right]$. Thus an automorphism $\varphi \in \text{Aut}(G_1 \times G_2)$ is determined by 4 group homomorphisms $\varphi_{ij} : G_j \rightarrow G_i$.

DEFINITION 2.1. We call $\varphi \in \text{Aut}(G_1 \times G_2)$, $G_1 \neq \{e\}$, $G_2 \neq \{e\}$, a diagonal automorphism if φ_{12} and φ_{21} are trivial homomorphisms.

DEFINITION 2.2. We call $(G_1 \times G_2) \rtimes_{\varphi} H$ a diagonal semidirect product if $\varphi(h)$ is a diagonal $G_1 \times G_2$ -automorphism for any $h \in H$. Explicitly, there are group homomorphisms $\varphi_{ii}(h) : G_i \rightarrow G_i$ such that $\varphi(h)(g_1, g_2) = (\varphi_{11}(h)(g_1), \varphi_{22}(h)(g_2))$.

REMARK 2.3. $\text{Ker}(\varphi) = \text{Ker}(\varphi_{11}) \cap \text{Ker}(\varphi_{22})$. Note that G_i may not be indecomposable as direct factors. Described encodings and diagonal semidirect products can be generalized to cases when the base groups splits into an arbitrary finite number of direct factors. Similar encodings can be used considering internal semidirect products.

2.1.2. Semidirect decompositions of diagonal semidirect products.

We present a proposition showing nonuniqueness of semidirect decomposition for diagonal semidirect products. Vaguely speaking, any direct factor of the base group which is an orbit of the initial twisting homomorphism can be moved to the extending group (nonnormal semidirect factor) to enlarge it. The new twisting homomorphism is such that the moved direct factor acts trivially on the remaining part of the base group.

PROPOSITION 2.4. Let $G = (N_1 \times N_2) \rtimes_{\varphi} H$ be a diagonal semidirect product, $\varphi(h)(g_1, g_2) = (\varphi_{11}(h)(g_1), \varphi_{22}(h)(g_2))$. Then the following statements hold.

1. $G \simeq N_1 \rtimes_{\Phi_{11}} (N_2 \rtimes_{\varphi_{22}} H)$, for some $\Phi_{11} \in \text{Hom}(N_2 \rtimes_{\varphi_{22}} H, \text{Aut}(N_2))$.

2. $\text{Ker}(\Phi_{11}) = \widetilde{N_2 \rtimes_{\varphi_{22}} H} \text{Ker}(\varphi_{11})$.

3. If $\varphi_{11}(h) = \text{id}_{N_1}$, for any $h \in H$, i.e. $\varphi(h)\left(\begin{bmatrix} g_1 \\ g_2 \end{bmatrix}\right) = \left[\begin{array}{c|c} g_1 & e \\ \hline e & \varphi_{22}(h)(g_2) \end{array}\right]$,

then

$$G \simeq N_1 \times (N_2 \rtimes_{\varphi_{22}} H).$$

Proof.

1. Consider $N_1 \rtimes_{\Phi_{11}} (N_2 \rtimes_{\varphi_{22}} H)$ where $\Phi_{11}(n_2, h) = \varphi_{11}(h)$. It is directly checked

that $\Phi_{11} \in \text{Hom}(N_2 \rtimes H, \text{Aut}(N_1))$. We will prove that

$$(N_1 \times N_2) \rtimes_{\varphi} H \simeq N_1 \rtimes_{\Phi_{11}} (N_2 \rtimes_{\varphi_{22}} H).$$

Define a bijective map $f : (N_1 \times N_2) \rtimes_{\varphi} H \rightarrow N_1 \rtimes_{\Phi_{11}} (N_2 \rtimes_{\varphi_{22}} H)$ by $f((n_1, n_2), h) = (n_1, (n_2, h))$, for all $n_i \in N_i$, $h \in H$. We prove that f is a group homomorphism.

Let $a, a' \in N_1$, $b, b' \in N_2$, $h, h' \in H$. We have that

$$((a, b), h) \cdot ((a', b'), h') = ((a, b)\varphi(h)(a', b'), hh') =$$

$$= ((a, b)(\varphi_{11}(h)(a'), \varphi_{22}(h)(b')), hh') = ((a\varphi_{11}(h)(a'), b\varphi_{22}(h)(b')), hh').$$

On the other hand, $(a, (b, h)) \cdot (a', (b', h')) = (a\Phi_{11}(b, h)(a'), (b, h) \cdot (b', h')) = (a\varphi_{11}(h)(a'), (b\varphi_{22}(h)(b'), hh'))$. We see that f is a group isomorphism.

2. $\text{Ker}(\Phi) = \{(n_2, h) | h \in \text{Ker}(\varphi_{11})\} = \widetilde{N_2} \widetilde{\text{Ker}(\varphi_{11})}$.

3. In notations given above, $\varphi_{11}(h) = id_{N_1}$ implies $\Phi_{11}(n_2, h) = id_{N_1}$, for any $n_2 \in N_2$, $h \in H$. Thus it is the direct product. \square

EXAMPLE 2.5. Let $G = (\mathbb{Z}_7 \times \mathbb{Z}_9) \rtimes_{\varphi} \mathbb{Z}_3$, where $\varphi(1)\left(\begin{bmatrix} g_1 \\ g_2 \end{bmatrix}\right) = \left[\begin{array}{c|c} g_1^2 & e \\ e & g_2^4 \end{array}\right]$. In additive notation this can be simplified as follows. $\varphi(1)\left(\begin{bmatrix} g_1 \\ g_2 \end{bmatrix}\right) = \left[\begin{array}{c|c} 2g_1 & 0 \\ 0 & 4g_2 \end{array}\right] = \left[\begin{array}{c|c} 2 & 0 \\ 0 & 4 \end{array}\right] \left[\begin{bmatrix} g_1 \\ g_2 \end{bmatrix}\right]$. G can be defined as the subgroup of Σ_{16} generated by three permutations:

- a) $(1, \dots, 7)$ (generating \mathbb{Z}_7),
- b) $(8, \dots, 16)$ (generating \mathbb{Z}_9) and
- c) $\underbrace{(1, 2, 4)(3, 6, 5)}_{\mathbb{Z}_7} \underbrace{(8, 11, 14)(9, 15, 12)}_{\mathbb{Z}_9}$ (generating action of \mathbb{Z}_3 on $\mathbb{Z}_7 \times \mathbb{Z}_9$).

We have that $G \simeq \mathbb{Z}_7 \rtimes (\mathbb{Z}_9 \rtimes_4 \mathbb{Z}_3) \simeq \mathbb{Z}_9 \rtimes (\mathbb{Z}_7 \rtimes_2 \mathbb{Z}_3)$.

2.2. Applications.

2.2.1. Generalized dihedral groups.

We remind the reader that an external semidirect product $D(A) = A \rtimes_{\varphi} \mathbb{Z}_2$ is called *generalized dihedral group* provided 1) A is abelian and 2) $\varphi(1)(g) = -g$ for any $g \in A$, in additive notation. We can also denote $D(A)$ by $A \rtimes_{-1} \mathbb{Z}_2$.

Using the known classification of finite abelian groups we can assume that $A = \bigoplus_{i=1}^n \mathbb{Z}_{m_i}$. We use linear algebra style encoding - we encode $(g_1, \dots, g_n) \in A$ as a column vector $\begin{bmatrix} g_1 \\ \dots \\ g_n \end{bmatrix}$. Notations introduced in section 2.1.1 are modified for additive group notation. The action of the twisting homomorphism is given by scalar or matrix multiplication: $\varphi(1)\left(\begin{bmatrix} g_1 \\ \dots \\ g_n \end{bmatrix}\right) = \left[\begin{array}{c|c|c} (-g_1) & 0 & 0 \\ \hline 0 & \dots & 0 \\ \hline 0 & 0 & (-g_n) \end{array}\right] = -\begin{bmatrix} g_1 \\ \dots \\ g_n \end{bmatrix} = (-\mathbf{E}_n) \cdot \begin{bmatrix} g_1 \\ \dots \\ g_n \end{bmatrix}$, where \mathbf{E}_n is the $n \times n$ identity matrix.

REMARK 2.6. *Generalized dihedral groups are diagonal semidirect products.*

PROPOSITION 2.7. *Let $A = \bigoplus_{i=1}^n \mathbb{Z}_{m_i}$, let $A = A_1 \oplus A_2$, where $A_1 = \bigoplus_{i=1}^{n_1} \mathbb{Z}_{m_i}$, $A_2 = \bigoplus_{i=n_1+1}^n \mathbb{Z}_{m_i}$. Then*

$$D(A) \simeq A_1 \rtimes (A_2 \rtimes_{-1} \mathbb{Z}_2) = A_1 \rtimes D(A_2) \simeq A_1 \rtimes D(A/A_1).$$

Proof. $D(A) = (A_1 \oplus A_2) \rtimes_{\varphi} \mathbb{Z}_2$, where $\varphi(1)(g) = -g$, for any $g \in A$. Thus $\varphi(g_1, g_2) = (-g_1, -g_2)$, for any $g_i \in G_i$. It follows that $D(A)$ is a diagonal semidirect product with respect to $A_1 \oplus A_2$ decomposition. According to Proposition 2.4 we have that $D(A) \simeq A_1 \rtimes_{\Phi_{11}} (A_2 \rtimes_{\varphi_{22}} \mathbb{Z}_2) = A_1 \rtimes D(A_2)$, where $\Phi_{11}(g_2, 1)(g_1) = \varphi_{11}(1)(g_1) = -g_1$. \square

EXAMPLE 2.8. *Let $G = D(\mathbb{Z}_3 \oplus \mathbb{Z}_5)$. G can be defined as a subgroup of Σ_8 generated by permutations $(1, 2, 3)$, $(4, 5, 6, 7, 8)$ and $(1, 2)(4, 7)(5, 6)$. Then $G \simeq \mathbb{Z}_3 \rtimes D_{2 \cdot 5} \simeq \mathbb{Z}_5 \rtimes D_{2 \cdot 3}$.*

2.2.2. Dihedral groups.

Classic dihedral groups are special cases of generalized dihedral groups when the base group is a cyclic group. We give a complete description of semidirect decompositions of D_{2n} using both Proposition 2.4 and ad hoc computations.

We use a classical presentation of dihedral groups:

$$D_{2n} = \langle a, x \mid a^n = e, x^2 = e, xax = a^{n-1} \rangle = \langle a \rangle \cup \langle a \rangle x.$$

We note that $D_2 \simeq \mathbb{Z}_2$ and $D_4 \simeq \mathbb{Z}_2 \times \mathbb{Z}_2$, in all other cases D_{2n} is nonabelian.

Subgroups. Let $n \in \mathbb{N}$, $n \geq 3$, $d \in \mathbb{N}$, $d \mid n$, $m = \frac{n}{d}$. It is known that D_{2n} has the following subgroups, see [2].

1. For each $m \in \mathbb{N}$ such that $m|n$ there is a subgroup

$$A_m = \langle a^{\frac{n}{m}} \rangle = \langle a^d \rangle = \{e, a^d, a^{2d}, \dots, a^{(m-1)d}\} \simeq \mathbb{Z}_m.$$

$A_m \leq D_{2n}$ for all m . The number of such subgroups is $d(n)$ (the number of natural n -divisors).

2. For each $m \in \mathbb{N}$ such that $m|n$ and each $r \in \mathbb{Z}_{\frac{n}{m}} = \mathbb{Z}_d$ there is a subgroup

$$B_{2m,r} = \langle a^{\frac{n}{m}}, a^r x \rangle = \langle a^d, a^r x \rangle = A_m \cup A_m(a^r x) \simeq D_{2m}.$$

The number of such subgroups is $\sigma(n)$.

If $2|n$ then $B_{n,r} \leq D_{2n}$. In all other cases, if $1 < m < n$ then $B_{2m,r} \not\leq D_{2n}$.

Classical decompositions. It known that $D_{2n} \simeq \mathbb{Z}_n \rtimes_{\varphi} \mathbb{Z}_2$ where the twisting homomorphism is $\varphi(1)(g) = -g$. In internal terms, $D_{2n} = A_n \rtimes B_{2,r}$, for all $r \in \mathbb{Z}_n$. If $2|n$ and $4 \nmid n$, then $D_{2n} \simeq D_n \times \mathbb{Z}_2$, or, in internal terms, $D_{2n} = B_{n,r} \times A_2$. where $r \in \mathbb{Z}_2$.

External semidirect decompositions of D_{2n} . Using Proposition 2.4 we get an exhaustive description of external semidirect decompositions of D_{2n} .

PROPOSITION 2.9.

1. $D_{2n} \simeq \mathbb{Z}_m \rtimes_{\varphi} D_{\frac{2n}{m}}$, for any $m \in \mathbb{N}$, $m|n$, such that $\text{GCD}(m, \frac{n}{m}) = 1$. φ is defined as follows: if $D_{\frac{2n}{m}} = \langle a, x | a^{\frac{2n}{m}} = e, x^2 = e, xax = a^{-1} \rangle$ then $\varphi(a)(1) = 1$ and $\varphi(x)(1) = -1$.
2. $D_{2n} \simeq D_n \rtimes_{\varphi} \mathbb{Z}_2$, if $n = 2^{\alpha}q$, $\alpha \in \mathbb{N}$. φ is defined as follows: if $D_n = \langle a, x | a^{\frac{n}{2}} = e, x^2 = e, xax = a^{-1} \rangle$ then $\varphi(1)(a) = a^{-1}$ and $\varphi(1)(x) = ax$.
3. If $2|n$ and $4 \nmid n$ then

$$D_{2n} \simeq D_n \times \mathbb{Z}_2.$$

4. There are no other nontrivial external semidirect decompositions of D_{2n} in the following sense. If $D_{2n} \simeq X \rtimes Y$, $|X| > 1$, $|Y| > 1$, then there are two possibilities:
 - a) $X = \mathbb{Z}_m$ and $Y = D_{\frac{2n}{m}}$, where $m|n$, $\text{GCD}(m, \frac{n}{m}) = 1$ or
 - b) $X = D_n$ and $Y = \mathbb{Z}_2$, if $2|n$.

Proof. Statements 1.,2.,3. are proved by exhibiting a suitable internal semidirect decomposition.

1. We use the primary decomposition theorem for cyclic groups: if $n = \prod_{i=1}^k p_i^{\alpha_i}$, then $\mathbb{Z}_n \simeq \bigoplus_{i=1}^k \mathbb{Z}_{p_i^{\alpha_i}}$. The statement follows from Proposition 2.4.

Alternatively, we prove the same statement using the information about D_{2n} -subgroups. We show that if $GCD(m, \frac{n}{m}) = 1$ then $D_{2n} = A_m \rtimes B_{\frac{2n}{m}, r}$.

We have that $A_m \leq D_{2n}$ and $|A_m| \cdot |B_{\frac{2n}{m}, r}| = 2n = |D_{2n}|$. $A_m \cap B_{\frac{2n}{m}} \leq \langle a^{\frac{n}{m}} \rangle$. Considering subgroups of $\langle a^{\frac{n}{m}} \rangle$ it follows from Lemma 2.10 that $A_m \cap B_{\frac{2n}{m}} = \{e\}$. Thus $D_{2n} = A_m \rtimes B_{\frac{2n}{m}, r} \simeq_{\varphi} \mathbb{Z}_m \rtimes_{\varphi} D_{\frac{2n}{m}}$. A direct computation shows that φ is as stated: $(a^m)a^d(a^{-m}) = a^d$, $(a^r x)a^d(a^r x) = a^{-d}$.

Note that if $2|n$ and $4 \nmid n$ then $A_2 \cap B_{n, r} = \{e\}$, $r \in \mathbb{Z}_2$, hence $D_{2n} = A_2 \times B_{n, r} \simeq \mathbb{Z}_2 \times D_n$. In this case there are no nontrivial semidirect decompositions of type $\mathbb{Z}_2 \rtimes D_n$.

2. This case is not covered by Proposition 2.4, we show directly that $D_{2n} = B_{n, 0} \rtimes B_{2, 1}$.

If $2|n$, then $B_{n, 0} \leq D_{2n}$. $|B_{n, 0}| \cdot |B_{2, 1}| = |D_{2n}|$. It can be checked that $B_{n, 0} \cap B_{2, 1} = \{e\}$: $B_{n, 0} = \langle a^2, x \rangle$, $B_{2, 1} = \langle ax \rangle$.

Thus $D_{2n} \simeq D_n \rtimes \mathbb{Z}_2$. A direct computation shows that φ is as stated: $(ax)a^2(ax) = a^{-2}$ (the generator a^2 gets inverted), $(ax)x(ax) = a^2x$ (the generator x gets multiplied by the other generator a^2).

3. Using Proposition 2.4 we see that $D_{2n} = D(\mathbb{Z}_n) = (\mathbb{Z}_2 \oplus \dots) \rtimes \mathbb{Z}_2 \simeq \mathbb{Z}_2 \times D(\mathbb{Z}_n/\mathbb{Z}_2) \simeq \mathbb{Z}_2 \times D_n$.

It can also be proved using the list of subgroups. We remind that $D_{2n} = B_{n, 0} \times A_2 \simeq D_n \times \mathbb{Z}_2$ for the following reasons. Both subgroups are normal. $|B_{n, 0}| \cdot |A_2| = |D_{2n}|$. $B_{n, 0} = \langle a^2, x \rangle$, $A_2 = \langle a^{\frac{n}{2}} \rangle$, $\frac{n}{2}$ is odd, therefore $B_{n, 0} \cap A_2 = \{e\}$.

4. Consider all possible internal semidirect decompositions of D_{2n} .

If $D_{2n} = X \rtimes Y$ then X must be a normal subgroup of D_{2n} therefore X must be A_m or $B_{n, r}$ with $2|n$.

If $X = A_m$ then Y must be $B_{m', r}$ in order to generate D_{2n} , with $m' = \frac{2n}{m}$. $A_m \cap B_{\frac{2n}{m}, r} = \{e\}$ iff $GCD(n, \frac{n}{m}) = 1$ by Lemma 2.10.

Let $X = B_{n, r}$ with $2|n$, $r \in \mathbb{Z}_2$. There are $n + 1$ subgroups of D_{2n} having order 2: $B_{2, r}$, $r \in \mathbb{Z}_n$ and $A_2 = \langle a^{\frac{n}{2}} \rangle$. For any n such that $2|n$ this gives a semidirect decomposition of type $D_n \rtimes \mathbb{Z}_2$. If $4 \nmid n$ then $A_2 \cap B_{n, r} = \{e\}$ which gives a direct decomposition $D_n \times \mathbb{Z}_2$. \square

LEMMA 2.10. *Let $n, a, b \in \mathbb{N}$, $a|n$, $b|n$. Then*

$$|\mathbb{Z}_a \cap \mathbb{Z}_b| > 1 \text{ iff } GCD(a, b) > 1.$$

Proof. As it was noted above, if $m|n$ then we identify \mathbb{Z}_m with the appropriate subgroup of \mathbb{Z}_n . Direct computations show that $GCD(a, b) = d$ iff $LCM(\frac{n}{a}, \frac{n}{b}) = \frac{n}{d}$.

Using the multiplicative notation of cyclic groups we can assume that $\mathbb{Z}_n = \langle z | z^n = e \rangle$, $\mathbb{Z}_a = \langle z^{\frac{n}{a}} \rangle$, $\mathbb{Z}_b = \langle z^{\frac{n}{b}} \rangle$. $|\mathbb{Z}_a \cap \mathbb{Z}_b| > 1$ iff $LCM(\frac{n}{a}, \frac{n}{b}) < n$ iff $GCD(a, b) > 1$. \square

REMARK 2.11. *In terms of prime factorization the condition $GCD(m, \frac{n}{m}) = 1$ is equivalent to the fact that m and $\frac{n}{m}$ are products of full prime powers of the prime factorization of n . Existence of many members of this family also follows from Schur-Zassenhaus theorem. If $m|n$ and $GCD(m, \frac{n}{m}) = 1$ then $GCD(|A_m|, |D_{2n}/A_m|) = 1$, $D_{2n}/A_m \simeq D_{\frac{2n}{m}}$ and, hence $D_{2n} \simeq A_m \rtimes D_{\frac{2n}{m}}$.*

REMARK 2.12. *Note that there are at most 2 external semidirect decompositions when n is a prime power:*

1. *if $n = p^\alpha$, p an odd prime, then there is only one (classical) external semidirect decomposition: $D_{2p^\alpha} \simeq \mathbb{Z}_{p^\alpha} \rtimes \mathbb{Z}_2$,*
2. *if $n = 2^\alpha$, $\alpha \geq 3$, then there are two external semidirect decompositions: $D_{2 \cdot 2^\alpha} \simeq \mathbb{Z}_{2^\alpha} \rtimes \mathbb{Z}_2 \simeq D_{2^\alpha} \rtimes \mathbb{Z}_2$.*

REMARK 2.13. *The image of the twisting homomorphism in each case of a proper semidirect product is isomorphic to \mathbb{Z}_2 .*

EXAMPLE 2.14.

External semidirect decompositions of $D_{2 \cdot 30}$:

$$\begin{aligned} D_{60} &\simeq \mathbb{Z}_{30} \rtimes \mathbb{Z}_2 \simeq \mathbb{Z}_6 \rtimes D_{10} \simeq \mathbb{Z}_{10} \rtimes D_6 \simeq \mathbb{Z}_{15} \rtimes D_4 \simeq \mathbb{Z}_3 \rtimes D_{20} \simeq \mathbb{Z}_5 \rtimes D_{12} \simeq \\ &\simeq D_{30} \rtimes \mathbb{Z}_2 \simeq D_{30} \times \mathbb{Z}_2. \end{aligned}$$

Internal semidirect decompositions of D_{2n} . We now describe all internal semidirect decompositions of D_{2n} .

PROPOSITION 2.15. *Let $n \in \mathbb{N}$.*

1. *If $m \in \mathbb{N}$, $m|n$, is such that $GCD(m, \frac{n}{m}) = 1$, then*

$$D_{2n} = A_m \rtimes B_{\frac{2n}{m}, r},$$

for all $r \in \mathbb{Z}_m$.

2. *If $n = 2^\alpha q$, $\alpha \in \mathbb{N}$, then*

$$D_{2n} = B_{n,0} \rtimes B_{2,r_1} = B_{n,1} \rtimes B_{2,r_0}$$

where $r_i \in \mathbb{Z}_n$, $r_i \equiv i \pmod{2}$.

3. If $2|n$ and $4 \nmid n$ then $D_{2n} = B_{n,0} \times A_2$ and $D_{2n} = B_{n,1} \times A_2$.
4. There are no other internal semidirect decompositions of D_{2n} .

Proof.

1. We look for internal semidirect decompositions of D_{2n} in form $A_m \rtimes B_{m',r}$. We must have $m' = \frac{2n}{m}$ and $r \in \mathbb{Z}_m$. $A_m \cap B_{\frac{2n}{m},r} = \{e\}$ iff $GCD(m, \frac{n}{m}) = 1$. Thus $D_{2n} = A_m \rtimes B_{\frac{2n}{m},r}$ for all m such that $GCD(m, \frac{n}{m}) = 1$ and all $r \in \mathbb{Z}_m$ are the only possible decompositions of this kind.

2. We look for internal semidirect decompositions of D_{2n} in form $B_{m,r} \rtimes B_{m',r'}$. We must have $B_{m,r} \trianglelefteq D_{2n}$ therefore $2|n$, $m = n$ and $r \in \mathbb{Z}_2$, thus we have two possible decomposition series: $B_{n,0} \rtimes B_{2,r'}$ and $B_{n,1} \rtimes B_{2,r''}$. To ensure trivial intersections of semidirect factors we must have $r' \equiv 1(mod 2)$ and $r'' \equiv 0(mod 2)$.

3. If $2|n$ and $4 \nmid n$ then $B_{n,0} \cap A_2 = B_{n,1} \cap A_2 = \{e\}$ where all subgroups are normal.

4. It follows from the previous arguments. \square

3. Conclusion. We have obtained a result showing possibility of various semidirect decompositions of a given semidirect product if the original twisting homomorphism is diagonal. This result may stimulate further interest in developing Krull-Remak-Schmidt theorem type results for semidirect and Zappa-Szep products.

As an application we have presented semidirect decompositions of generalized dihedral groups and all possible external and internal semidirect decompositions of classical dihedral groups. Apart from semidirect decompositions guaranteed by the general proposition, for D_{2n} there are additional decompositions of external type $D_n \rtimes \mathbb{Z}_2$ if $2|n$.

Acknowledgement. Computations were performed using the computational algebra system MAGMA, see Bosma et al. [1].

REFERENCES

- [1] Bosma W, Cannon J, and Playoust C (1997), The Magma algebra system. I. The user language, J. Symbolic Comput., 24, pp. 235-265.
- [2] Conrad K (2015) Dihedral groups II <http://www.math.uconn.edu/~kconrad/blurbs/grouptheory/dihedral2.pdf>
- [3] Hillar C, Rhea D (2007) Automorphisms of finite abelian groups American Mathematical Monthly, 114, no.10, pp. 917-923.
- [4] Robinson D (1996) A course in the theory of groups. Springer-Verlag, New York.
- [5] Rotman J (1995) An introduction to the theory of groups. Springer-Verlag, New York.